

Contents

Introduction.....	2
Overview.....	2
Storage and Stability.....	2
Binding Capacity.....	2
Kit Contents.....	3
Before Starting.....	4
E.Z.N.A.™ Yeast RNA Spin Protocol.....	4
On-Membrane DNase I Digestion Protocol.....	6
Quantization and Storage of RNA.....	7
RNA Quality.....	7
Troubleshooting Guide.....	8

Revised October 2006

Introduction

The E.Z.N.A.™ Yeast RNA Kit allows convenient isolation of high-quality total RNA from a wide variety of yeast species. Up to 2×10^7 log-phase cultured yeast cells can be processed. The system combines the reversible nucleic acid-binding properties of HiBind® matrix with the speed and versatility of spin column technology to yield approximately 30 µg of RNA, with an A_{260}/A_{280} ratio of 1.7-1.9. Purified RNA is suitable for downstream applications such as RT-PCR, DD-PCR, and hybridization techniques. There are no organic extractions, thus reducing plastic waste and hands-on time to allow multiple samples to be processed in parallel.

Overview

If working with RNA for the first time, please read this booklet to become familiar with the procedure. Yeast cells are grown to log-phase and spheroblasts are subsequently prepared using Buffer SE and lyticase. Following lysis, binding conditions are adjusted and the sample is applied to a HiBind® RNA spin-column. Two rapid wash steps remove trace salt and protein contaminants, and RNA is eluted in DEPC-treated water. Purified RNA can be directly used in downstream applications without need for further purification.

Storage and Stability

All components of the E.Z.N.A.™ Yeast RNA Kit are stable for at least 24 months from date of purchase when stored at 22°C-25°C. During shipment or storage in cool ambient conditions, precipitates may form in Buffer YRL. These precipitates should be dissolved by warming the solution at 37°C and gently shaking its container.

Binding Capacity

Each HiBind RNA Mini Column can bind up to 100 µg Total RNA. Use no more than 3 ml log yeast culture is not recommended.

Kit Contents

Product Number	R6870-00	R6870-01	R6870-02
Purification times	5 Preps	50 Preps	200 Preps
HiBind® RNA Mini column	5	50	200
Glass Beads	270mg	2.7 g	12 g
2 ml Collection Tubes	15	150	600
Buffer SE	15 ml	120 ml	420 ml
Buffer YRL	5 ml	20 ml	80 ml
RNA Wash Buffer I	5 ml	45 ml	180 ml
RNA Wash Buffer II	5 ml	12 ml	2 x 25 ml
Lyticase	1400 units	14000 units	56,000 units
DEPC-water	1 ml	20 ml	50 ml
User Manual	1	1	1

*Buffer YRL contains a chaotropic salt. Use gloves and protective eye wear when handling this solution.

Before Starting

- Whenever working with RNA, always wear latex gloves to minimize RNase contamination. Use only clean, RNase-free disposable plastic pipette tips when using the supplied reagents.
- During the procedure work carefully but quickly.
- Dissolve the lyticase with water to 4000 U/ml and store at -20°C.

R6870-00	Dissolve with 350µl water
R6870-01	Dissolve with 3.5ml water
R6870-02	Dissolve with 35ml water

- Under cool ambient conditions, crystals may form in Buffer YRL. This is normal and the bottle should be warmed to redissolve the salt.
- 2-mercaptoethanol (β-mercaptoethanol) is key in denaturing RNases and must be added to an aliquot of Buffer YRL before use. Add 20-µl of 2-mercaptoethanol per 1 ml of Buffer YRL. This mixture can be stored for 1 month at room temperature.

- RNA Wash Buffer II is supplied as a concentrate and must be diluted with absolute ethanol as follows and **stored at room temperature**.

R6870-00	Add 20 ml absolute (96%-100%) ethanol
R6870-01	Add 48 ml absolute (96%-100%) ethanol
R6870-02	Add 100 ml absolute (96%-100%) ethanol

E.Z.N.A.™ Yeast RNA Kit Spin Protocol

This protocol uses enzymatic lysis to prepare spheroblasts. Freshly prepare lyticase/SE Buffer by mixing 63µl of lyticase (4000 U/ml) with 2ml SE Buffer and add 20µl of 2-mercaptoethanol before use.

Materials to Be Provided by User

- Microcentrifuge and nuclease-free 2.0 and 1.5 ml tubes.
- Swinging-bucket centrifuge and sterile 15 ml tubes.
- Waterbath equilibrated to 30°C.
- 70% ethanol - do not use other alcohols.
- Absolute ethanol (96-100%)
- 2-Mercaptoethanol

Sample size

While the HiBind® RNA mini columns can bind up to 100 µg RNA, for effective purification, use no more than 5×10^7 log-phase yeast cells. For *Scerevisiae* grown in YPD, an OD_{600} of 1.0 corresponds to approximately 2×10^7 cells per ml.

1. **Collect no more than 5×10^7 yeast cells** in a 15 ml tube by centrifuging for 5 min at 1000 x g at 4°C. **NOTE:** Use only freshly harvested cells for preparation of spheroblasts.
2. **Aspirate and discard supernatant completely and resuspend cells in 2 ml SE /Lyticase mixture.** Incubate at 30°C for 30 min, inverting the tube once every 10 mins. Spheroblasts must be handled gently.

Note: Depend on the yeast strain, the incubation time, amount of lyticase and composition of SE Buffer may vary.
3. **Pellet spheroblasts by centrifuging 5 min at 400 x g at room temperature.** Carefully aspirate and discard supernatant. Incomplete

removal of supernatant will prevent complete lysis of spheroblasts in the next step.

4. Add 350 µl of Buffer YRL/2-mercaptoethanol and 50mg **glass beads** into the tube. Vortex at max speed for 5 minutes to lyse and homogenize the sample. **If bead mill is available, lyse the cell with bead mill at top speed until the cells are completely lysed.**
5. Centrifuge at 13,000 x g for 3 min. Transfer the supernatant into a new centrifuge tube.
6. Add an equal volume 70% ethanol to the lysate and mix thoroughly by pipetting or vortexing at max speed for 15 seconds. A white precipitate may form upon addition of ethanol; it will not interfere with the procedure. **Do not centrifuge the tube.**
7. Apply the entire sample (around 700 µl) to a HiBind® RNA Mini column assembled in a 2 ml collection tube (supplied). Centrifuge at $\geq 10,000$ xg for 30 seconds **at room temperature**. Discard flow-through and re-use collection tube in Step 8 or Step 9.

Note: This is the point at which **optional** on-membrance DNase I digestion should begin. If DNase I digestion is desired, skip Step 8 and proceed to Step 9. If DNase I digestion is not needed, proceed with Step 8.

8. Wash the sample by adding 500µl of RNA Wash Buffer I to the column. Centrifuge at 10,000 xg for 15 seconds at room temperature. Discard the flow-through and re-use the collection tube. (This is starting point for On-Membrance DNase Digestion protocol, See "Optional On-Membrance DNase Digestion Protocol" on page 7).
9. **Place column in the same 2 ml collection tube**, and add 500 µl RNA Wash Buffer II diluted with ethanol. Centrifuge at 10,000 xg for 15 seconds and discard flow-through. Re-use the collection tube in next step.

Note: Wash Buffer II Concentrate must be diluted with absolute ethanol before use. Refer to Page 4 or to label on bottle for directions.

10. Wash sample with a second 500 µl of Wash Buffer II and centrifuge and discard flow-through as in preceding step. Using the same collection tube, centrifuge the spin cartridge at 10,000 x g for **1 min at full speed** to completely dry the HiBind® matrix.
11. **Elution of RNA.** Transfer the column to a clean 1.5 ml microfuge tube (not supplied with kit) and elute the RNA with 50-100 µl of DEPC-treated water (supplied with kit). Make sure to add water directly onto column matrix. Centrifuge 1 min at maximum speed. A second elution may be necessary if the expected yield of RNA is greater than 50 µg. Alternatively, RNA may be eluted with a greater volume of water. While additional elutions increase total

RNA yield, the concentration will be lowered since more than 80% of RNA is recovered with the first elution. Pre-heating the water to 70°C before adding to column and incubating column 5 min at room temperature before centrifugation may increase yields.

On-Membrance DNase I Digestion Protocol (Optional)

Since HiBind® RNA resin and spin-column technology actually removes most DNA without the DNase treatment, this DNase I digestion step is not necessary for most downstream applications. However, certain sensitive RNA applications might require further DNA removal. (See DNase I, Cat # E1091 for detailed information.)

1. For each HiBind® RNA column, have the following DNase I digestion reaction mixture prepared in advance as follows:

OBI DNase I Digestion Buffer	73.5 µl
RNase-free DNase I (20 Kunitz units/µl)	1.5 µl
Total volume	75 µl

Note:

- A. **DNase I is very sensitive, and is thus subject to physical denaturation, so do not vortex this DNase I mixture. Mix gently by inverting the tube. Prepare the fresh DNase I digestion mixture before RNA isolation.**
 - B. **OBI DNase I digestion buffer is supplied with the OBI RNase-free DNase set.**
 - C. **Standard DNase buffers may not be compatible with the Omega Bio-Tek kit for on-membrance DNase digestion.**
2. Dry column by spinning an additional 30 seconds, then pipet 75 µl of the DNase I digestion reaction mixture directly onto the surface of the HiBind® RNA membrane in each column. Make sure to pipet the DNase I digestion mixture directly onto the membrane. DNase I digestion might not be complete if some of the mixture sticks to the wall or to the O-ring of the HiBind® RNA column.
 3. Incubate at room temperature (25-30°C) for 15 minutes.
 4. Place column in a clean 2 ml collection tube and add 350 µl RNA Wash Buffer I. Allow wash buffer to soak column for at least 5 minutes before proceeding. Centrifuge at 10,000 x g for 15 seconds at room temperature. Discard flow-through and re-use the collection tube in next

step.

Quantitation and Storage of RNA

To determine the concentration and purity of RNA, measure absorbance at 260 nm and 280 nm in a spectrophotometer. 1 O.D. unit measured at 260 nm corresponds to 40 µg of RNA per ml. DEPC-water is slightly acidic and can dramatically lower absorbance values. We suggest that you dilute the sample in a buffered solution (TE) for spectrophotometric analysis. The ratio of A_{260}/A_{280} of pure nucleic acids is 2.0, while for pure protein it is approximately 0.6. A ratio of 1.8-2.0 corresponds to 90%-100% pure nucleic acid. Store RNA samples at -70°C in water. Under such conditions RNA prepared with the E.Z.N.A.® system is stable for more than a year.

RNA Quality

It is highly recommended that RNA quality be determined prior to all analyses. The quality of RNA can best be assessed by denaturing agarose gel electrophoresis and ethidium bromide staining. Two sharp bands should appear on the gel. These are the 28S and 18S ribosomal RNA bands. If these band smear towards lower molecular weight RNAs, then the RNA has undergone major degradation during preparation, handling, or storage.

Troubleshooting Guide

Problem	Cause	Suggestion
Little or no RNA eluted	RNA remains on the column	<ul style="list-style-type: none"> Repeat elution. Pre-heat DEPC-water to 70° C prior to elution. Incubate column for 5 min with water prior to centrifugation.
	Column is overloaded	<ul style="list-style-type: none"> Reduce quantity of starting material.
Clogged column	Incomplete spheroblasting	<ul style="list-style-type: none"> Extend incubation with Lyticase. Increase Lyticase to 200 units per 10⁷ cells. Reduce amount of starting material
Degraded RNA	Source	<ul style="list-style-type: none"> Use only freshly harvested cells. Do not store cells prior to extraction unless they are lysed with Buffer YRL first. Follow protocol closely, and work quickly.
	RNase contamination	<ul style="list-style-type: none"> Ensure not to introduce RNase during the procedure. Check buffers for RNase contamination.
Problem in downstream applications	Salt carry-over during elution	<ul style="list-style-type: none"> Ensure Wash Buffer II Concentrate has been diluted with 4 volumes of 100% ethanol as indicated on bottle. Wash Buffer II must be stored and used at room temperature. Repeat wash with Wash Buffer II.
DNA contamination		<ul style="list-style-type: none"> Perform the optional Dnase I digestion step or Digest with RNase-free DNase after elution and inactivate at 75°C for 5 min.
Low Abs ratios	RNA diluted in acidic buffer or DEPC-water	<ul style="list-style-type: none"> DEPC-treated water is acidic and can dramatically lower Abs₂₆₀ values. Use TE buffer at pH 8.0 to dilute RNA prior to spectrophotometric analysis.